

Investigating age related changes in taste and affects on sensory perceptions of oral nutritional supplements

Article

Accepted Version

Kennedy, O., Law, C., Methven, L., Mottram, D. and Gosney, M. (2010) Investigating age related changes in taste and affects on sensory perceptions of oral nutritional supplements. *Age and Ageing*, 39 (6). pp. 733-738. ISSN 0002-0729 doi: <https://doi.org/10.1093/ageing/afq104> Available at <https://centaur.reading.ac.uk/8136/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1093/ageing/afq104>

Publisher: Oxford University Press

Publisher statement: This is a pre-copy-editing, author-produced PDF of an article accepted for publication in [insert journal title] following peer review. The definitive publisher-authenticated version: Kennedy, O., Law, C., Methven, L., Mottram, D. and Gosney, M. (2010) Investigating age related changes in taste and affects on sensory perceptions of oral nutritional supplements. *Age and Ageing* , 39 (6). pp. 733-738. <http://ageing.oxfordjournals.org/content/39/6/733.abstract>

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Investigating age related changes in taste and affects on sensory perceptions of oral nutritional supplements.

Background: Sip feeds are oral nutritional supplements (ONS) that are commonly prescribed to malnourished patients to improve their nutritional and clinical status. However, ONS are poorly consumed and frequently wasted, with sweetness being identified as one of the factors leading to patients' dislike of ONS.

Objectives: To investigate if age affects sweetness thresholds and if this impacts upon perceived sweetness intensity, hedonic (sweetness and overall) and ranked preference of ONS products.

Design: prospective, observational.

Subjects: Thirty six young adults (18-33 years) and 48 healthy older adults (63-85 years).

Setting: Dept. of Food & Nutritional Sciences and the Clinical Health Sciences at the University of Reading.

Methods:

Detection and recognition threshold levels, basic taste identification and 'just about right' level of sweetness were examined. Three ONS (chocolate, vanilla, strawberry) and sucrose solutions were evaluated for hedonic sweetness, overall hedonic liking, sweetness intensity and rank preference.

Results

Significant differences were found in both sweetness detection and recognition thresholds ($P=0.0001$) between young and older adults, with older adults more likely to incorrectly identify the taste ($P=0.0001$). Despite the deterioration in sweetness sensitivity among the older adults, there were no significant differences found in sweetness intensity perceived for the ONS products presented ($P>0.05$) when compared to the young adults. However, across both groups sweetness intensity was found to be correlated with overall product dislike across all flavour variants tested ($R=0.398$, $p=0.0001$).

Conclusions: Sweetness appears to be one of many factors contributing to the dislike of ONS. Manufacturers are encouraged to reconsider the formulations of these products so that beneficial effects of ONS can be delivered in a more palatable and acceptable form and wastage reduced.

Keywords: oral nutritional supplement, taste, acceptability, preference

Introduction

In the United Kingdom the population is ageing with 16% of the population aged 65 and over [1]. Healthy ageing is essential to maintain a high quality of life and is defined by World Health Organisations (WHO) as the state of complete physical, mental and social well being [2]. However, malnutrition among older adults diminishes quality of life by contributing to serious illness, decreased functional capability and altered self-perception of health and chronic disability [3]. The prevalence of malnutrition in older adults is widespread across the U.K. and varies geographically, with higher prevalence in the North (19.4%) than the South (11.2%) of England [4]. Recent statistics from National Institute for Health and Clinical Excellence [5] suggested that more than 10% of over 65's in the general population are at medium or high risk of malnutrition, and that this figure rises to 60% in the hospital setting. The 2008 British Association for Parenteral and Enteral Nutrition (BAPEN) Nutrition screening survey found that one in three adults admitted to hospitals was malnourished and that those aged 65 plus, had 40% greater risk of malnutrition than those <65 years [6].

Many authors, including early work by McWhirter and Pennington [7], have suggested that the earlier nutritional intervention is started, the greater the clinical benefit to patients. This will in turn lower the total cost of treating malnutrition to the

UK National Health Service (NHS), estimated at £7.4 billion [8] and to the UK; £13 million per annum [9]. Nutritional intervention therapies such as dietary supplements are commonly used to treat malnutrition, and can be given enterally or parenterally [10]. Oral supplements such as ONS ready to drink supplements available in cartons, cans, bottles and cups, can be provided on prescription to patients and used in addition to normal food consumption to increase nutrient intake. The typical nutritional content of an ONS is 1.5 Kcal per ml and consists of 14-20% protein, 25-35% fat, 50%-60% carbohydrates; the vitamin and mineral content generally supplies a proportion of the Recommended Nutrient Intake (RNI) in a 200ml pack [11]. Much previous research and reviews have examined the prevention/treatment of malnutrition using products such as these. [10, 13, 14].

However, it is reported that the use of ONS is associated with high wastage, therefore, the benefit of ONS cannot be delivered if they are not consumed. Gosney [13] investigated the palatability and consumption of ONS and the reasons for wastage. Wastage in four wards (96 patients) within a 24 hour period was as much as 63% and this wastage was further extrapolated to a net loss of £18,924 on ONS per year our elderly care wards in the 1990's. Sweetness was given as one of the reasons leading to dislike of these products.

Aim of this study

This aim of the study was to investigate if a difference in sweetness perception as indicated by threshold and detection levels exist between young and older adults and to examine if this was related to differences, if any, in the perception of sensory attributes, acceptability and hedonic liking of ONS. Our null hypothesis is that differences in sweetness thresholds between young and older adults are so small that they will not relate to the perception of sensory attributes, acceptability and hedonic liking of ONS.

Methods and materials.

The study was approved by the Research and Ethics Committee at the University of Reading, UK prior to recruiting the study participants.

Recruitment of young and older adult participants

The young adults were recruited by advertising through email and posters from students and staff at the University of Reading. Older adults were recruited via postal communication with those on the older adults' database held by Clinical Health Sciences at the University of Reading. All self selected respondents who met the inclusion criteria were invited to participate in the study; therefore a convenience sampling strategy was employed. Thirty-six young adults age 18-33 (mean age 23)

Age related sensory changes & ONS

and forty-eight healthy older adults age 63-85 (mean age 74.4) participated in this study.

Sensory tests

Stimuli for sweetness threshold tests

The series of sucrose solutions for testing were prepared according to British Standard ISO 3972:1991[15]. Commercially available spring water (Harrogate Spa) was used throughout the study. All solutions were prepared within 24-hours prior to each test in order to retain freshness and to prevent separation.

Stimuli for preference tests

Ensure Plus a commercially available nutritional sip feed was chosen for this study as it was identified to be the most often prescribed brand within the elderly care wards at the local NHS Trust Three ONS flavours; vanilla, strawberry and chocolate, were chosen as these were the most commonly prescribed product variants. All had identical nutritional values and sweetness levels.

Procedures

All the sensory sessions for the young adults took part in a dedicated sensory laboratory, and the sensory tests for the older adults took place at the Clinical Health

Age related sensory changes & ONS

Sciences, both located at the University of Reading. Participants in both groups completed two sets of tests on each of three separate occasions; a series of threshold tests and a series of sensory attribute intensity, hedonic liking and rank preference tests. Each sample in each test was coded with a randomised 3 digit code and samples in the second set of tests were presented in randomised order. Water and crackers (Carr's brand) were provided as palate cleansing tools to minimise carry over effects.

Threshold tests

Participants compared each of nine solutions with the reference sample(water) and identified the point where they could detect a difference (detection threshold) and the point where they could identify the taste (recognition threshold) which they had perceived. Detection threshold is a sensory term used to describe the concentration (or level) at which an individual (or group of individuals) can detect the existence of a signal from the background noise. To calculate a threshold the participant is given a series of increasing stimuli (in this case concentrations of sucrose in water), the detection threshold is calculated as the geometric mean between two samples, the highest concentration at which the participant could not detect the signal and the lowest concentration at which they could. Both thresholds enable the study to identify the sensitivity of sweetness in each participant. Participants were also required to choose the Just About Right (JAR) concentration of the presented solutions.

Sensory attribute & preference tests

Volunteers were presented with three flavours of the sip feeds and a sucrose solution containing the equivalent sugar content (28.8g in 500ml/5.76g in 100ml), presented using a random balanced order design. The sucrose solution acted as a control,

Comment [OBK1]: Suggest that this goes into supplementary information as will take the words to below 2500

enabling the investigation of how flavour type impacts on sweetness perception. Sweetness liking/hedonic and the overall product liking of the samples using a 7 point hedonic scale (like extremely (1) to dislike extremely (7)). Following this, participants were asked to rank products in order of overall preference (1 most preferred to 3 least preferred). Sweetness intensity was measured on a 100mm line scale with descriptors 'not at all sweet' (0mm) to 'extremely sweet' (100mm). Participants were required to rate how much (quantitatively) they liked the sweetness intensity.

Statistical analysis:

Data analysis was performed using Statistical Package for the Social Sciences (SPSS) (Chicago, Illinois, Version 15.0). As no differences were found in intra-individual results between test days ($P>0.05$), means of the triplicate test results were calculated for each participant and were used throughout for analysis (to compare between age groups and genders). A two way analysis of variance (ANOVA) was used to examine the impact of age and gender on the variables measured with post hoc analysis, the multiple pair-wise comparisons test used to compare samples was Fishers least significant difference (LSD) ($P<0.05$).

Results

No differences were found in intra-individual results between test days ($P>0.05$), therefore means of the triplicate test results were calculated for each participant and were used throughout for analysis (to compare between age groups and genders).

Threshold tests-sweetness sensitivity differences in young and older adults.

A significant difference in detection threshold ($P=0.0001$), recognition threshold ($P=0.0001$) and Just About Right (JAR) sweetness level ($P=0.03$) was found between the young and older adults. The older adults were significantly less sensitive to sweetness than the young adults. A significant difference in the older adults compared to the younger group ability to identify the taste was found ($P=0.0001$ data not shown). Gender did not affect any of the parameters measured.

Table 1. Mean detection and recognition thresholds, Just About Right (JAR) sweetness levels by age and gender.

	Young adults	Older	Young males	Young	Older	Older
		adults		females	Males	females
Thresholds (g/L)	Mean	Mean	Mean	Mean	Mean	Mean
Detection	3.75 ^a	5.52 ^b	3.45 ^a	3.87 ^a	5.10 ^b	5.78 ^b
Recognition	5.94 ^a	6.89 ^b	5.77 ^a	6.00 ^a	6.79 ^b	6.95 ^b
JAR	6.34 ^a	6.87 ^b	5.92 ^a	6.51 ^b	6.76 ^b	6.94 ^b

Mean values within the same row with different superscripts are significantly different as determined by Fishers least significant difference (LSD) (P<0.05).

Product hedonic liking tests

In overall hedonic liking (both young and older adults), chocolate was the most liked and the control sucrose solution was the least liked sample tested. This data is also mirrored by the rank preference data where the chocolate variant was ranked more preferred for both young and older adults.

Table 2. Mean product overall hedonic liking and sweetness intensity for all products across all groups.

	Vanilla	Strawberry	Chocolate	Sucrose
	Mean	Mean	Mean	Mean
Overall hedonics\$	3.77 ^a	3.91 ^a	3.60 ^a	4.44 ^b
Sweetness intensity*	64.5 ^a	68.5 ^b	54.2 ^c	75.4 ^d

*Mean values within the same row with different superscripts are significantly different as determined by Fishers least significant difference (LSD) (P<0.05) \$ 7 point hedonic scale; *100mm line scale*

Table 3. Overall hedonic liking of products rating by age group and gender.

	Young adults	Older	Young	Young	Older	Older
		adults	males	females	Males	females
Hedonic liking*	Mean	Mean	Mean	Mean	Mean	Mean
Vanilla	3.71 ^a	3.81 ^a	3.05 ^a	3.97 ^a	2.81 ^a	4.4 ^b
Strawberry	3.88 ^a	3.92 ^a	3.35 ^a	4.08 ^a	2.98 ^a	4.49 ^b
Chocolate	3.41 ^a	3.74 ^a	3.22 ^a	3.49 ^a	3.45 ^a	3.91 ^a
Sucrose	4.07 ^a	4.71 ^a	4.85 ^a	3.76 ^b	4.31 ^a	4.97 ^a

*7 point hedonic scale

*Mean values within the same row with different superscripts are significantly different as determined**by Fishers least significant difference (LSD) (P<0.05)*

Please see the tables 4 & 5 Appendix 2 in the supplementary data on the journal

website

<http://www.ageing.oxfordjournals.org/>*Product sweetness intensity*

The sweetness intensity of the sucrose solution perceived by the older adults was

Age related sensory changes & ONS

significantly lower than ratings from the young group ($P=0.03$), and young males perceived the solutions as significantly sweeter than all other groups ($P<0.05$). No differences were found between other flavour variants in terms of sweetness intensity ratings by either age or gender.

Despite the identical sweetness level across all samples, the sweetness intensity perceived by both groups (young and older adults) were significantly different for each product variant, with the chocolate flavour sweetness intensity perceived as significantly lower than all other variants and the sucrose solution as the highest sweetness intensity ($P<0.05$). This suggests that there is a strong taste and/ aroma interaction in perceptions of sweetness intensity. It is possible that coca would increase bitterness which may have suppressed the sweetness.

Relationship between sweetness intensity and hedonic measures

An inverse relationship was found for overall and individual product variants sweetness intensity and both hedonic liking of the product sweetness ($R=0.399$, $P=0.001$) and overall product characteristics ($R=0.398$, $P=0.001$). It may therefore be inferred that as the products perceived sweetness intensity increased, so too did product dislike.

Discussion and conclusion.

Significant differences were found in both detection and recognition thresholds and the correct identification of the basic taste of sweetness between young and older adults, which reflects the findings of other researchers [16, 17, 18] who have reported that taste sensitivity decreases with age. It has been suggested that differences in taste recognition, such as those found in this study may be due to a time lag in the turnover of taste receptor cells [19] as work has shown that there is no decrease in the number of taste buds with age which earlier research had postulated [20,21].

Although previous work has demonstrated aged related changes in taste acuity, few studies have explored the interaction of this with real food systems such as ONS used in the current study. Work carried out in the Netherlands found older adults perceived sweet ingredients in chocolate drinks and vanilla waffles as less intense than the young [22, 23]. However no differences in perceived sweetness of dairy products [24] or custards [25] has been found.

In the current study, dislike of ONS was found, the degree of which varied across flavours, genders and age groups. The chocolate flavour ONS was found to be the most liked, and was perceived as less sweet than other product variants presented in this study. In general, all older adults who had significantly higher sweetness thresholds rated the ONS more negatively for liking (both sweetness and overall liking of the samples) although they perceived the sweetness intensity of the products

244 to be less intensive than the young adults. This infers that other product sensory
245 attributes in addition to sweetness are contributing to the dislike of these products.
246 Indeed, it has been suggested by others that the incongruence found between sensory
247 and hedonic modalities is due to different processing pathways for these in the brain
248 [26, 27).

249 It had been assumed in the past that age related sensory losses may lead to
250 modifications of food pleasantness and food choice [28], however others suggest a
251 habituation process to the decreased perception exists which may offer a
252 compensatory mechanism to prevent decreases in food liking [29]. This may in part
253 explain why significant differences in taste acuity found in the current study did not
254 translate into significant differences in liking across the age groups.

255 In Gosney's study [13], 38% of participants disliked the sweetness of ONS (not
256 flavour specific) which reflects the results of the current study where 27% of the older
257 adults disliking the sweetness, whereas in young adults this was more accepted as
258 only 6% disliked the sweetness. However in this study, 25% disliked that taste, 19%
259 the texture and a further 19% commented that they felt sick or bloated post
260 consumption. Compliance was extremely low at only 37%. Although we did not find
261 that sweetness acuity directly related to liking of these products, we did find that
262 sweetness intensity is one of the factors leading to dislike of these products.

Age related sensory changes & ONS

263 Even though the difference in results shown in table 3, may appear small and some
264 may question their clinical relevance, however in terms of consumer acceptability,
265 products with a value of <5 , are unlikely to be available on the open market, these low
266 scores indicate poor liking of the products, but show that the degree of this was
267 moderated by flavour. In addition, the difference between liking scores of over 1 point
268 (on a 7 point scale) for some of the flavour variants between older males and females
269 is of interest and concern. Liking scores of <3 , as a mean for older males, would
270 certainly be expected to lead to product rejection. It is also interesting to note, that in
271 general males across both age groups scored the products less favourably than females,
272 again increasing the likelihood of their rejection.

273 We acknowledge that we used a convenience sample of young and older adults, and
274 therefore our results may not be generalisable, however we feel that they are
275 important in pointing out some directions in which research could be focused in terms
276 of effective treatment of malnutrition. It has long been known that ONS are wasted
277 with the common reason given that consumers/patients dislike the taste, our
278 research has sought to address what aspects taste are disliked, and examine if this is
279 related to sweetness thresholds and sweetness and flavour of the products. The
280 research which we have presented in this paper has used a multidisciplinary approach
281 involving sensory scientists, flavour chemists, dietitians, nutritionists and clinicians to

Age related sensory changes & ONS

try and understand some of the ‘taste’ issues surrounding poor consumption and we are currently using these finding to develop solutions to improve ONS acceptability.

We would suggest that more research is needed in this area as results of this study indicate that ONS may need to be reformulated depending on target group (age and gender) to increase acceptance and increase consumption. Reformulation will be especially beneficial amongst malnourished older adults but also within all groups where these products are used, so that required nutrients could be delivered in a more palatable form, maximising the nutritional effects of ONS and in turn reducing wastage.

Acknowledgements

Abbott Nutrition are thanked for the supply of ONS used in this study.

Key points

- Young and older adults possess significantly different sweetness threshold levels.
- As perceived products sweetness increased, liking of the products decreased.
- Chocolate flavour ONS was the most preferred product tested.

Conflict of Interest:

None declared.

References

[1]ONS. Key population and vital statistics-2007. 2009; available at <http://statistics.gov.uk> . Accessed September 02, 2009.

[2]Chen CCH, Schilling LS, Lyder CH. A concept analysis of malnutrition in the elderly. *J Adv Nurs* 2001; 36: 131-142.

[3]Millen BE. Preventive nutrition services for aging populations. In *Malnutrition in the Elderly* (Seiler W.O and Strahelin H.B. eds), Springer, New York, 1999; 121-132.

[4]Elia M and Stratton RJ. Geographic inequalities in malnutrition prevalence in the elderly across England. *Proc Nutrition Soc* 2005, 64: 12A.

[5]NICE. National Institute for Health and Clinical Excellence. *New NICE guideline will help tackle the problem of malnutrition in NHS. 2006*; Available at: <http://www.nice.org.uk>. Accessed September 02, 2009.

[6]BAPEN.Nutrition screening survey in the UK in 2008. 2009; Available at: <http://www.bapen.org.uk>. Accessed September 02, 2009.

[7]McWhirter JP and Pennington CR (1994) Incidence and recognition of malnutrition in hospital. *BMJ* 1994; 303: 945- 948.

[8]Elia M, Pang F, Russell C, Green C. and Stratton RJ. *The Cost of Malnutrition in*

- 321 *the UK and the Economic Case for Oral Nutritional Supplement (ONS) in Adults.*
- 322 Redditch, Worcs: BAPEN, 2005.
- 323 [9] BAPEN. *Combating Malnutrition: Recommendations for Action*. 2009; available
- 324 at: <http://www.bapen.org.uk>. Accessed September 02, 2009.
- 325 [10]Stratton RJ, Green CJ and Elia M. Disease-Related Malnutrition: an evidence
- 326 based approach to treatment. In: *Consequences of Disease-related Malnutrition*.
- 327 CABI Publishing, Wallingford 2003; 4: 113- 155.
- 328 [11]Ralph A. Human Nutrition and Dietetics. In : *Dietary Reference Value 10th edition*.
- 329 (Garrow J.S., James W.P.T and Ralph A. eds), Churchill Livingstone, London, 2000;
- 330 829-863.
- 331 [12] Koretz RL, Avenell A, Lipman TO, Braunschweig CL and Milne AC. Does
- 332 enteral nutrition affect clinical outcome? A systematic review of the randomised trials.
- 333 Am J Gastroenter 2007; 102; 412-429.
- 334 [13]Gosney M. Are we wasting our money on food supplements in elder care wards? J
- 335 Adv Nurs 2003; 43: 275-280.
- 336 [14] Milne AC, Potter J, Vivanti, A and Avenell, A. Protein and energy
- 337 supplementation in elderly people at risk from malnutrition. Cochrane Database Syst
- 338 Rev. 2009, April 15:CD003288.
- 339 [15] British Standard Method for sensory analysis of food. Part 7: Investigating
- 340 sensitivity of taste. BS 5929-7: 1992, ISO 3972:1991[16] Mojet J, Christ-Hazelhof E,

- 341 Heidema J. Taste perception with age generic or specific losses in threshold sensitivity
342 to the five basic tastes? *Chem Senses* 2001; 26: 845-60.
- 343 [17] Fukunaga A, Uematsu H, Sugimoto K. Influences of aging on taste perception
344 and oral somatic sensation. *J Gerontol A* 2005; 60: 109–113.
- 345 [18] Wardwell L, Chapman-Novakofski K, Brewer MS. Effects of age, gender and
346 chronic obstructive pulmonary disease on taste acuity. *Int J Food Sci Nutr.* 2009; 19:
347 1-14.
- 348 [19] Sugimoto K, Iseki H. Morphological characteristics of the taste bud in aged mice.
349 *Jpn J Taste Smell Res* 1994; 1: 234-236.
- 350 [20] Arvidson K. Location and variation in number of taste buds in human fungiform
351 papillae. *J Dent Res.* 1979; 87: 435-442.
- 352 [21] Bradley RM, Stedman HM, Mistretta CM. Age does not affect numbers of taste
353 buds and papillae in adult rhesus monkeys. *Anat Rec.* 1985; 212: 246-24
- 354 [22] Mojet J, Heidema J, Christ-Hazelhof E. Taste perception with age: generic or
355 specific losses in supra-threshold intensities of five taste qualities? *Chem Senses* 2003;
356 28: 397–413.
- 357 [23] Kremer S, Mojet J, Kroeze JHA. Differences in perception of sweet and savoury
358 waffles between elderly and young subjects. *Food Qual Pref* 2007; 18: 106–116.

- 359 [24]De Graaf C, Polet P, van Staveren W. Sensory perception and pleasantness of food
360 flavours in elderly subjects. *J Gerontol* 1994; 49: 93–99.
- 361 [25] Kremer S, Bult JH, Mojet J, Kroeze JH. Food perception with age and its
362 relationship to pleasantness. *Chem Senses* 2007; 32: 591-602.
- 363 [26] De Araujo IET, Rolls ET, Kringelbach ML, McGlone F, Phillips N. Taste-olfactory
364 convergence, and the representation of the pleasantness of flavour, in the human brain. *Eur J*
365 *Neurosci* 2003; 18: 2059–2068.
- 366 [27] Sedwards TV. Dual separate pathways for sensory and hedonic aspects of taste.
367 *Brain Res Bull* 2004; 62: 271–283.
- 368 [28] Mattes RD. The chemical senses and nutrition in aging: challenging old
369 assumptions. *J Am Diet Assoc* 2002; 102: 192–196.
- 370 [29] Wysocki CJ, Pelchat ML. The effects of aging on the human sense of smell and
371 its relationship to food choice. *Crit Rev Food Sci Nutr* 1993; 33: 63–82.
- 372
- 373
- 374

375 Supplementary data

376 **Table 4.** Flavour rank preference data according to age group.

	Most liked (came first in preference ranking)	Least liked
Young adults	Chocolate	Strawberry
Older adults	Chocolate	Vanilla

377

378 **Table 5.** Mean product sweetness intensity perception rating by age group and gender.

	Young adults	Older	Young	Young	Older	Older
		adults	males	females	Males	females
Sweetness intensity*	Mean	Mean	Mean	Mean	Mean	Mean
Vanilla	6.45 ^a	6.45 ^a	5.94 ^a	6.64 ^a	6.07 ^a	6.68 ^a
Strawberry	6.86 ^a	6.75 ^a	6.62 ^a	6.96 ^a	6.52 ^a	6.89 ^a
Chocolate	5.55 ^a	5.32 ^a	5.16 ^a	5.70 ^a	5.24 ^a	5.37 ^a
Sucrose	8.03 ^a	7.17 ^b	9.02 ^a	7.63 ^b	7.05 ^b	7.25 ^b

*7 point hedonic scale

379 *Mean values within the same row with different superscripts are significantly different as*

380 *determined by Fishers least significant difference (LSD) (P<0.05).*

381

Age related sensory changes & ONS

